Carex Backii Boott (C. durifolia Bailey).

Glyceria nervata Trin., var. stricta Scribn.

Panicum Werneri Scribn.

" tennesseense Ashe.

Phegopteris hexagonoptera Fée. A large colony on a shaded hillside, July, 1905.

Aspidium spinulosum Swartz (typical). On a rocky hill in partial shade, the plants evidently increasing.

South Chesterville, Maine.

The Dwarf Mistletoe on the Southeastern Coast of Maine.

— The recent articles calling attention to Arceuthobium pusillum at different points in Maine prompt this note, supplementing the series of Maine records.

The plant first came to my attention during July, 1902, at West Jonesport, occurring rarely on Black Spruce, on the main land. Late in July, 1904, at Cross Island, off the mouth of Machias Bay, the plant was found in great profusion, again on Black Spruce. Here it infested patches of limited extent, with larger patches free from infection.

On Matinicus Island, off the coast of Maine, on August 4th, 1907, this plant was found in greater abundance and size than I have seen it previously. It infested the small, exceedingly dense or "scrubby" White Spruces about the Northwest Head.

I have often looked, vainly, for it about Casco Bay, yet we are now warranted in considering that it has been shown to occur throughout the northern, central, and eastern portions of the State. The regularity of its distribution is another question, deserving of many reports.

— ARTHUR H. NORTON, Portland, Me.

```
The following records are to be found in Rhodora.
Vol. ii. 2 (1900): von Schrenk — Monhegan Island, Boothbay, Linekin. 10 (1900): Fernald — Fort Kent, Upsala. 221 (1900): Arthur — Isle au Haut.
Vol. v. 49 (1903): Harvey — Mt. Katahdin.
Vol. viii, 168 (1906): Chamberlain — Pleasant Ridge.
Vol. xi. 28 (1907): Murdoch — Rangeley. 75 (1907): Rand — Mt. Desert Island. 124 (1907): Chamberlain — Norway — Casco.
[Arceuthobium is also found in Alton and Georgetown — Eds.]
```

Vol. 9, No. 105, including pages 149-196 was issued 13 September, 1907.